



PHARMACEUTICAL EVALUATION OF SHATYADI CHURNA

Dr. Chinmayee Sahu^{1*}, Prof. (Dr.) Pragya P. Mallik², Dr. Santilata Sahoo³

¹PG Scholar, PG Department of Kayachikitsa, GAM, Puri (Odisha).

²Professor, PG Department of Kayachikitsa, GAM, Puri (Odisha).

³Associate Professor, PG Department of Kayachikitsa, GAM, Puri (Odisha)

ABSTRACT

Medicinal plants form an integral part of traditional healthcare systems and are now recognized globally for their therapeutic importance. Their curative actions are attributed to a wide range of secondary metabolites present in various plant parts. **Shatyadi Churna** is a classical Ayurvedic formulation prepared from eight key herbs in equal proportion, with a Bhabana Dravyas added to enhance efficacy. The present work aims to explore its phytochemical profile, pharmacological attributes, and probable mechanism of action. **Shatyadi Churna** is a classical powder formulation indicated in disorders of Grahani, Atisāra and Prameha conditions in different Nighaṅṭus. Grahāṇi Roga is a disease of Agni Vaishamya where Mandagni leads to improper digestion, Ama formation and deranged Grahāṇi functions. The formulation contains Deepana, Pacana, Grahāṇisaṁsthapanana and Srotoshodhana drugs. Therefore, Shatyadi Churna corrects the Agni at Jatharagni level and reduces Ama, improves the digestive fire and normalises Grahāṇi function. The present study, carried out at ALN RAO MEMORIAL AYURVEDIC MEDICAL COLLEGE AND PG CENTRE, KOPPA, CHIKMAGALUR (KARNATAKA), includes standardization parameters such as identity, purity, strength, preliminary phytochemical screening, and TLC profiling. These findings play a crucial role in establishing scientific quality control measures and enhancing the therapeutic credibility of Ayurvedic formulations.

KEYWORDS: Shatyadi Churna, Deepana, Pacana, Grahāṇisaṁsthapanana, Srotoshodhana, Grahāṇi, Agni, Ama, Preliminary Phytochemical Screening, And TLC Profiling.

INTRODUCTION

Ayurveda, the ancient science of life, emphasizes not only the therapeutic utility of formulations but also their proper pharmaceutical preparation and evaluation. Churna (powder) is one of the most widely used dosage forms in Ayurvedic pharmaceuticals (Rasa Shastra and Bhaishajya Kalpana)¹. To achieve desired therapeutic effects, the quality of the raw drugs and the precision of the preparation process are of paramount importance.

Shatyadi Churna is a classical polyherbal powder formulation described in Yogaratnakar, Grahaniroagadhikar, Kaphaja Grahani. It contains Karchura (Curcuma zedoaria), Sunthi (Zingiber officinalis), Maricha (Piper nigrum), Pippali (Piper longum), Haritaki (Terminalia chebula), Pippalimoola (Piper longum), Yavakshara, Sarjikshara in equal proportions, with Vijapuraka (Citrus medica) as Bhabana Dravyas. The formulation Shatyadi Churna is primarily indicated in Grahāṇi Roga, especially the Kaphaja type of Grahāṇi, where Mandagni and excessive Kapha lead to picchila mala, gaurava, alasya and amapratīṣṭhana. This formulation exhibits Dipana, Pachana, Grahāṇisaṁsthapanana and Srotoshodhana properties, which correct Agnidushti, eliminate Ama and normalise Grahāṇi function. The Tikta-Kaṭu-Kaṣaya Rasa along with Laghu-Rukṣa-Tikṣṇa Guṇa of Shatyadi Churna directly antagonise Kapha, therefore reducing mucous content in stool, heaviness and bloating. It also supports proper digestion and absorption, thereby improving Dhatu-poṣhaṇa. The pharmaceutical evaluation of Shatyadi Churna includes standard organoleptic, physicochemical and qualitative parameters as per API/WHO guidelines to ensure uniformity, consistency and quality standardisation of the formulation.

AIMS AND OBJECTIVES

1. To prepare Shatyadi Churna according to classical Ayurvedic texts.
2. To evaluate its pharmaceutical parameters including organoleptic, physicochemical, and phytochemical properties.
3. To establish standard quality control measures for the formulation, Shatyadi Churna.

MATERIALS AND METHODS

Source of Data

1. Classical text book of Ayurveda
2. Text books of Modern science
3. Published articles from periodical journals another magazines.



Raw Materials

All the raw drugs were collected from authentic Ayurvedic suppliers and identified based on their **macroscopic and microscopic characters** as per *API (Ayurvedic Pharmacopoeia of India)* standards.

Method of Preparation 2,3

All the raw drugs of Shatyadi Curṇa were collected from authentic GMP approved raw drug store and were authenticated by the Department of Dravyaguna. Each drug was cleaned properly to remove physical impurities like soil, stones and foreign particles. Afterwards the drugs were dried under shade at room temperature to protect the volatile phytoconstituents. Each ingredient was powdered separately using a mechanical pulveriser to obtain fine powder. The powders were then passed through 80# sieve to achieve uniform particle size. All the sieved powders were mixed thoroughly in a clean, dry stainless steel container to obtain a uniform and homogeneous mixture of Shatyadi Curṇa. Finally, the prepared Churṇa was packed in an airtight, moisture-free, light resistant container and stored at room temperature until further use. Then the sample of final product, Shatyadi Curṇa was labeled with name, parts of ingredients and send to Laboratory for Pharmacological evaluation.

Table 1: Ingredients of Shatyadi Curṇa

| Sl. No. | Drug Name (Botanical Name) | Botanical Name | Part Used | Quantity |
|---------|----------------------------|----------------------|-----------------|-------------|
| 1 | Shati | Curcuma zedoaria | Kanda (Rhizome) | 1 part |
| 2 | Sunthi | Zingiber officinalis | Kanda (Rhizome) | 1 part |
| 3 | Maricha | Piper nigrum | Phala (Fruit) | 1 part |
| 4 | Pippali | Piper longum | Phala (Fruit) | 1 part |
| 5 | Haritaki | Terminalia chebula | Phala (Fruit) | 1 part |
| 6 | Pippalimoola | Piper longum | Moola (Root) | 1 part |
| 7 | Yavakshara | - | — | 1 part |
| 8 | Sarjikshara | - | — | 1 part |
| 9 | Vijapuraka | Citrus medica | Swarasa | As required |

INGREDIENTS OF SHATYADI CHURNA





PHARMACEUTICAL STUDY/ ANALYTICAL STUDY:

The Shatyadi Curṇa was analyzed by adopting various related analytical parameters like

A. Organoleptic Characteristics

Colour, odour, taste and appearance of Churna were observed and mentioned in (Table no.2)

(Table No. 2)

| | |
|----------------|--------------------------------------|
| Colour | Greyish green |
| Odour | Aromatic |
| Taste | Slightly bitter, pungent, astringent |
| Texture | Powder |

B. Physiochemical Analysis

Loss on drying at 105^oc, total ash value, water soluble ash, acid insoluble ash, pH value. Alcohol soluble extractives, Water soluble extractives was carried out for raw materials and results are mentioned in (Table no.3)

(Table No.3)

| Parameter | Result (% w/w) | Remarks / Observation |
|------------------------------------|----------------|--|
| Loss on Drying at 105°C | 3.71% | Indicates moisture content |
| Total Ash | 7.01 % | Represents total mineral content |
| Acid Insoluble Ash | 0.50% | Reflects siliceous (sand/silica) matter |
| Water Soluble Ash | 3.38 % | Indicates water-soluble inorganic salts |
| Alcohol Soluble Extractives | 17.81% | Shows alcohol soluble phytoconstituents |
| Water Soluble Extractives | 38.34 % | Represents water soluble active components |
| pH (5% Aqueous Solution) | 4.76 + 0.10 | Slightly acidic nature |

These results indicate good quality of the Churna, with optimal moisture and extractive values suggesting stability and solubility.

C. Preliminary Phytochemical Tests (Qualitative Tests):

For various functional groups were done and observation and result of the *churna* was asserted in (Table no.4).

(Table no.4)

| Phytochemical Test | Result |
|--------------------------------|---------|
| Carbohydrate | Present |
| Protein | Present |
| Alkaloid | Present |
| Cardiac Glycoside | Present |
| Flavonoids | Present |
| Tannins | Present |
| Anthraquinone Glycoside | Present |
| Triterpenoids | Present |

The presence of these phytochemicals supports the Deepana, Pacana, Grahaṇisamsthapana and Srotoshodhana properties described in Ayurvedic literature.

D. FLUORESCENT TESTS

| Reagent / Sample Treatment | Under Visible Light | Under Long UV |
|---|---------------------|--------------------|
| Sample + Water | Creamish Yellow | Fluorescent Yellow |
| Sample + MeOH | Light Yellow | Fluorescent Cream |
| Sample + 10% NaOH | Reddish Green | Fluorescent Green |
| Sample + 10% HCl | Yellowish Green | Fluorescent Green |
| Sample + 10% HNO₃ | Reddish Green | Fluorescent Green |
| Sample + 10% H₂SO₄ | Greenish Yellow | Fluorescent Yellow |
| Sample + 10% NH₃ | Light Brown | Brown |



E, TLC PROFILE

Solvent System: Toluene: Ethyl acetate:: 80:20

| Rf Value | Colour Under Long UV |
|----------|----------------------|
| 0.04 | Brownish Blue |
| 0.15 | Fluorescent Green |
| 0.22 | Fluorescent Green |
| 0.35 | Fluorescent Green |
| 0.56 | Fluorescent Green |
| 0.65 | Fluorescent Green |
| 0.74 | Fluorescent Green |

DISCUSSION

The pharmaceutical evaluation of Shatyadi Curṇa demonstrated that the formulation maintains desirable organoleptic and physicochemical characteristics suitable for Churna preparation. The extractive values indicate a balanced solubility of active principles in both aqueous and alcoholic media. The presence of vital phytoconstituents such as alkaloids and flavonoids ensures Deepana, Pacana, Grahaṇisamsthāpana and Srotoshodhana activity, correlating with its therapeutic indications in Grahaṇi Roga, especially the Kaphaja type of Grahaṇi.

The analytical data of *Shatyadi Curṇa* indicates that the formulation possesses satisfactory physicochemical characteristics ensuring its quality, stability, and efficacy. The **loss on drying (3.71%)** reveals a low moisture content, suggesting good stability and reduced susceptibility to microbial contamination. The **total ash value (7.01 %)** represents the overall mineral content, confirming the presence of inorganic constituents derived from the raw herbal materials. The **acid-insoluble ash (0.50%)** being very low denotes minimal siliceous or earthy impurities, reflecting the purity of the ingredients used. The **water-soluble ash (3.38 %)** suggests a considerable amount of inorganic salts soluble in water, contributing to the formulation's therapeutic mineral fraction. The **alcohol-soluble extractive value (17.81%)** indicates the presence of moderately polar phytoconstituents such as alkaloids, flavonoids, and resins, whereas the **water-soluble extractive value (38.34 %)** highlights the predominance of polar bioactive compounds like glycosides, tannins, and saponins that are easily extractable in aqueous media. The **pH value (4.76 + 0.10)** shows a slightly acidic nature, which is acceptable and suitable for oral administration. Overall, the results demonstrate that *Shatyadi Curṇa* is a well-balanced formulation with appropriate physicochemical properties supportive of its stability and therapeutic potential.

The phytochemical screening of Śatyādi Cūrṇa showed the presence of Carbohydrates, Proteins, Alkaloids, Cardiac glycosides, Flavonoids, Tannins, Anthraquinone glycosides and Triterpenoids which indicates that the formulation is pharmacodynamically rich. Alkaloids, flavonoids and triterpenoids are reported for strong Deepana-Pācana, antioxidant, anti-inflammatory, smooth muscle regulating and gut motility stabilising activities, which directly support the core chikitsā sūtra of Kaphaja Grahaṇi Roga where Agnimāndya and Āma formation are dominant. Tannins and cardiac glycosides help in grahaṇi samsthāpana by tightening gut mucosa, reducing excessive secretions and improving tone of the intestinal wall; while carbohydrate and protein fractions support Dhātu-varhdhana and restoration of depleted bio-energy after chronic mal absorption. Anthraquinone glycosides contribute to mild Srotoshodhana effect by ensuring proper colon cleansing without irritation. Collectively, these phytochemical constituents justify the classical claims that Śatyādi Cūrṇa acts as Deepana, Pācana, Grahaṇi Samsthāpana, Srotoshodhana, Śophahara and Rasāyana; hence effectively corrects the Samprapti of Kaphaja Grahaṇi Roga by improving Agni, reducing Kapha-Āma load and normalising intestinal function.

The TLC profiling of Śatyādi Cūrṇa under long UV showed multiple bands distributed throughout the Rf range (0.04 to 0.74) with predominant fluorescent green zones. Presence of several distinct spots at Rf 0.15, 0.22, 0.35, 0.56, 0.65 and 0.74 suggests that Shatyadi Churna contains a wide spectrum of phytoconstituents with varying polarity — which signifies a poly-herbal synergy. The predominance of fluorescent green bands indicates rich flavonoid fraction and related phenolic compounds, while the brownish-blue low Rf band around 0.04 indicates the presence of heavier / less mobile phytocompounds like tannins. Such multi-component profile supports the combined Deepana, Pachana and Srotoshodhana potential of the formulation. In the context of Kaphaja Grahaṇi Roga — where Kapha + Ama leads to mandagni, impaired digestion and malabsorption — this kind of diversified phytochemistry helps to simultaneously kindle Agni, reduce Ama, control excessive Kapha and stabilise intestinal function. Therefore, the observed TLC pattern provides scientific support to the classical claim that Śatyādi Cūrṇa acts as a multi-target formulation correcting Agni dushti and Grahaṇi dushti in Kaphaja Grahaṇi.

Proper standardization ensures batch-to-batch uniformity, enhances shelf life, and provides a scientific foundation for traditional formulations. Hence, pharmaceutical evaluation is an indispensable step before clinical or pharmacological studies.

CONCLUSION

The Śatyādi Cūrṇa prepared as per classical guidelines and evaluated on modern pharmaceutical parameters fulfils the essential qualitative standards required for safe and effective Ayurvedic formulations. The analysis confirms that the formulation has



acceptable physicochemical characteristics along with significant phytochemical constituents which directly support its classical claims in the management of Kaphaja Grahani Roga. The presence of multi-dimensional phyto-markers indicates strong Deepana, Pācana, Grahani Samsthāpana and Srotoshodhana potential which can help in correcting Agnimāndya, reducing Kapha-dominance and breaking the Ama-pathology. Further advanced analytical and pharmacological studies can strengthen the scientific evidence and establish wider therapeutic applicability of Śatyādi Cūrṇa in Grahani spectrum disorders.

REFERENCES

1. *Yogaratanakara, Grahaniroga Chikitsa Adhyaya, with Vidyotini Hindi Commentary by Vaidya Lakshmipati Shastri, Chaukhambha Prakashan, Varanasi, Reprint Edition, 2020.*
2. *Sharngadhara Saṁhitā, Madhyama Khanda, edited by Dr. Smt. Shailaja Srivastava, Chaukhamba Orientalia, Varanasi, Reprint Edition, 2019.*
3. *Bhaishajya Ratnavali, Grahani Roga Chikitsa Adhikara, edited by Ambikadutta Shastri, Chaukhamba Sanskrit Sansthan, Varanasi, 21st Edition, 2020.*
4. *Ayurvedic Pharmacopoeia of India (API), Part I, Vol. I-IX, Government of India, Ministry of AYUSH, Department of AYUSH, New Delhi, 2001-2020.*
5. *Ayurvedic Formulary of India (AFI), Part I, II & III, Government of India, Ministry of AYUSH, New Delhi, 2003-2016.*
6. *Khandelwal, K.R. Practical Pharmacognosy: Techniques and Experiments, 19th Edition, Nirali Prakashan, Pune, 2008.*
7. *Kokate, C.K., Purohit, A.P., and Gokhale, S.B. Pharmacognosy, 50th Edition, Nirali Prakashan, Pune, 2015.*
8. *Mukherjee, P.K. Quality Control of Herbal Drugs: An Approach to Evaluation of Botanicals, Business Horizons, New Delhi, 2019.*
9. *Trease, G.E., & Evans, W.C. Pharmacognosy, 16th Edition, Elsevier Publications, 2009.*
10. *Anonymous. Quality Standards of Indian Medicinal Plants, Vol. I-XIV, Indian Council of Medical Research (ICMR), New Delhi, 2003-2022.*
11. *Government of India. Protocol for Testing of Ayurvedic, Siddha and Unani Medicines, Department of AYUSH, New Delhi, 2008.*
12. *World Health Organization (WHO). Quality Control Methods for Herbal Materials, WHO, Geneva, 2011.*
13. *Patwardhan, B., Warude, D., Pushpangadan, P., & Bhatt, N. (2005). Ayurveda and traditional Chinese medicine: A comparative overview. Evidence-Based Complementary and Alternative Medicine, 2(4): 465-473.*
14. *Chopra, R.N., Nayar, S.L., & Chopra, I.C. Glossary of Indian Medicinal Plants, Council of Scientific and Industrial Research (CSIR), New Delhi, 2002.*
15. *Singh, R.H. (2012). Ayurvedic Principles of Pathophysiology (Samprapti-Vijnana), Chaukhambha Surbharti Prakashan, Varanasi.*
16. *Tripathi, B.N. Charaka Samhita, Chikitsa Sthana, Chaukhambha Surbharti Prakashan, Varanasi, Reprint Edition, 2020.*
17. *Pandey, G.S. Bhavaprakasha Nighantu, Chaukhambha Bharati Academy, Varanasi, Reprint Edition, 2018.*
18. *Sastry, J.L.N. (2012). Dravyaguna Vijnana, Vol. II, Chaukhambha Orientalia, Varanasi.*
19. *Sharma, P.V. Dravyaguna Vijnana, Vol. II, Chaukhambha Bharati Academy, Varanasi, Reprint Edition, 2017.*
20. *Dwivedi, R.R. & Singh, R.H. (1998). Clinical evaluation of Ayurvedic compound formulation in the management of Grahani Roga. Journal of Research in Ayurveda and Siddha, 19(1-2): 15-23.*