



IMPACT OF ELEVATED CO₂ ON *in vitro* PROPAGATION OF IMPORTANT MEDICINAL PLANT *Nyctanthes arbor-tristis* L.

Pushkar Choudhary¹, Roshni Adiyecha²

¹Department of Botany, Centre of Education, I.I.T.E., Gandhinagar, India

²Department of Botany, Centre of Education, I.I.T.E., Gandhinagar, India

ABSTRACT

Nyctanthes arbor-tristis L. is an important ornamental as well as medicinal plant known for its flowers. The axillary nodes of this plant are cultured on Murashige and Skoog medium, which results in the direct organogenesis. The M.S. medium is supplemented with 3 mg/L Indole-3-acetic acid. The *in-vitro* regenerated shoots are sub-cultured on the MS medium supplemented with IAA and BAP and subjected to elevated level of CO₂ viz 0.5% and 2% in airtight CO₂ chamber. The comparative analysis is performed on various physical and biochemical parameters after an 8-week treatment at elevated CO₂. The results of the present study show that the enhanced CO₂ level proves beneficial for *in vitro* growth of *Nyctanthes arbor-tristis* L.

KEYWORDS:- *Nyctanthes arbor-tristis* L., Elevated CO₂, M.S. Medium, Direct Organogenesis

INTRODUCTION

Nyctanthes arbor-tristis L., commonly referred to as Night-flowering Jasmine or Parijat, is a plant renowned for its significant medicinal, ornamental, and cultural value. This versatile plant has been utilized extensively in traditional medicine, with every part of the plant, including its leaves, roots, stems, bark, and seeds, demonstrating therapeutic properties. (Himanshi Rawat *et al.*, 2021 ; Jaspal Singh *et al.*, 2021.). The leaf juice is commonly used to aid digestion, act as an antidote to reptile venom, and serve as a tonic, laxative, diaphoretic (promoting sweating), and diuretic (increasing urine output). Additionally, the leaves are employed in the treatment of an enlarged spleen, highlighting their role in addressing specific organ-related conditions.

Traditionally, powdered dried leaves are known to alleviate malaria symptoms and rheumatic joint pain, reflecting their importance in treating both infectious diseases and inflammatory conditions. (Ashokkumar, K., Dharshini, M., Janani, T. *et al.* ; 2024) Beyond these uses, the plant exhibits a wide array of pharmacological activities, including antihistaminic effects (to manage allergic reactions) and central nervous system (CNS) actions such as hypnotic (sleep-inducing), tranquilizing (calming), and anesthetic properties. Its pharmacological spectrum further includes analgesic (pain-relieving), anti-inflammatory, antipyretic (fever-reducing), antiulcer, anti-amoebic, anti-helminthic (against parasitic worms), anti-trypanosomal (against trypanosome parasites), antidepressant, antiviral, and immunomodulatory (immune-regulating) activities.

Phytochemical studies of *Nyctanthes arbor-tristis* L. have revealed the presence of a wide range of bioactive compounds, including phyosterols, phenolics, tannins, flavonoids, glycosides, and saponins. (Mansi , Rajhans, Sanjukta *et al.*, 2021; Samir Panda *et al.*; 2024) Glycosides and alkaloids are particularly abundant in the plant, with iridoid glycosides and phenylpropanoid glycosides being the most frequently identified glycosides. Among the alkaloids, nyctanthine is

found to be the most prominent, playing a significant role in the plant's therapeutic efficacy.

The seeds of the plant are rich in iridoid compounds and arbortristins (A, B, D, and E), which have demonstrated immunomodulatory properties and antileishmanial activity (effective against *Leishmania* parasites). The leaves contain a diverse array of bioactive compounds, such as 6,7-di-O-benzoylnystantoin, 6-O-trans-cinnamoyl-6 β -hydroxyloganin, 7-O-trans-cinnamoyl-6 β -hydroxyloganin, and desrhamnosylverbascoside, which are known for their potent anti-inflammatory and antipyretic effects. The stem is characterized by the presence of naringenin-4-O- α -glucopyranoside and 1- β -xylopyranoside, which contribute to the plant's overall medicinal value. (Champa Rani, Sunaina Chawla *et al.*; 2012)

Environmental changes, particularly the rising levels of carbon dioxide (CO₂) due to anthropogenic activities, have forced plants like *Nyctanthes arbor-tristis* L. to adapt by altering their secondary metabolism. Elevated CO₂ levels are known to stimulate the production of secondary metabolites, which are often associated with enhanced pharmacological activities. Studies involving *In vitro* cultivation of *Nyctanthes arbor-tristis* L. Under elevated CO₂ conditions have provided insights into these adaptive changes, offering valuable knowledge about how such environmental factors influence the plant's phytochemistry and potential medicinal applications.

MATERIAL AND METHODS

Young axillary buds of *Nyctanthes arbor-tristis* L. which was collected from the campus of IITE, Gandhinagar, were used as explants. The explants were surface sterilized with 0.1% HgCl₂ for 2 minutes, followed by washing with sterilized distilled water, and then sterilized with 1% sodium hypochlorite for 2 minutes. After sterilization, the explants were washed with distilled water 5-6 times and then inoculated onto MS (Murashige and Skoog) media, supplemented with 3 mg/L Indole-3-acetic acid (IAA).



Direct Organogenesis

Axillary buds of approximately 1 cm² in length were inoculated onto MS media containing 3% sucrose and 0.8% Agar-agar. The medium was supplemented with 3 mg/L IAA.

Once the shoots were regenerated and the axillary buds reached sufficient length, they were sub-cultured onto MS media with 2% sucrose, and supplemented with 3 mg/L IAA and 1 mg/L BAP (6-benzylaminopurine). The inoculated culture bottles were placed in an airtight CO₂ chamber with CO₂ levels maintained at 0.5% to 2%.

The pH of the medium was adjusted to 5.8 and autoclaved at 121°C (15 psi) for 20 minutes. The cultures were incubated at 25°C under a 16-hour photoperiod.

RESULT AND DISCUSSION

The current study investigated the *in vitro* propagation of *Nyctanthes arbor-tristis* L through direct organogenesis, focusing on the role of varying carbon dioxide (CO₂) concentrations in influencing growth, physiological responses, and biochemical composition. Axillary nodes of approximately 1 cm² length were inoculated on MS medium supplemented with 3 mg/L indole-3-acetic acid (IAA). Within 10 days, initial shoot formation was observed, indicating that the explants were responsive to the culture medium and hormonal treatment. By the 20th day, multiple shoots were observed, each producing an average of 3±0.8 leaves. This robust growth suggests that the chosen medium and hormonal conditions effectively supported organogenesis. However, unhealthy explants showed reduced growth and organogenesis, underscoring the importance of explant health in achieving optimal results.

After 25 days of culture, the shoots were sub cultured and transferred to a fresh MS medium containing 3 mg/L IAA and 20 g/L sucrose and placed in airtight chambers with controlled CO₂ concentrations of 0%, 0.5%, and 2.5%. Organogenesis was observed in all treatments by the 12th day, but significant differences in growth were noted based on the CO₂ levels. Plants cultured under 2.5% CO₂ demonstrated the most vigorous growth, producing an average of 6±0.1 leaves per shoot, which was significantly higher compared to the 0.5% CO₂ treatment, where shoots produced only 5±0.3 leaves on average. Notably, no shoot or leaf formation was observed in the absence of CO₂ (0% treatment), highlighting the essential role of CO₂ in plant development and organogenesis.

Micropropagation can also be conducted on sugar-free media under photoautotrophic conditions, utilizing elevated CO₂ as a carbon source. This method has been demonstrated to lower the risk of microbial contamination (Kozai, 1991; Pospisilova *et al.*, 1999; Deng and Donnelly, 1993; Yoon *et al.*, 2009). However, in the current study, when CO₂ was used as the sole carbon source (without sucrose supplementation), leaf explants survived only for 6–8 days before deteriorating. Similar outcomes were reported in photoautotrophic cultures of *tobacco* (Solarova *et al.*, 1989; Ticha, 1996), *Wrightia tomentosa* (Vyas and Purohit, 2003), and *Chlorophytum*


borivilianum (Joshi *et al.*, 2009), where plantlets failed to sustain growth in sucrose-free media (Solarova and Pospisilova, 1997). In contrast, potato plantlets were able to grow in sugar-free media, although media containing sucrose resulted in superior growth (Rahman and Alsadon, 2007). Additionally, shoot cultures of *Achras zapota* cultivated on sucrose-containing media under CO₂ free conditions exhibited a significant decline in all growth parameters.

Biochemical analyses revealed clear trends in nutrient and metabolite accumulation. Plants grown under 2.5% CO₂ exhibited the highest phosphorus content (2.85±1.04mg/g of leaf), **followed by the control plants, while those grown under 0.5% CO₂ showed the lowest phosphorus levels (1.65mg/g of leaf)**. Similar trends were observed for reducing sugar and protein content, with plants at 2.5% CO₂ recording the highest levels (38.5±2.04 mg/g and 198±4.02mg/g of leaf respectively), **followed by the control plants, and plants at 0.5% CO₂ having the lowest values (20.34±1.09mg/g and 106±2.08mg/g of leaf respectively)**. These results suggest that elevated CO₂ levels enhance nutrient assimilation and promote the synthesis of critical biomolecules required for plant growth.


Physiological parameters such as stomatal frequency and chlorophyll content also varied significantly with CO₂ levels. Plants grown under 2.5% CO₂ exhibited the highest stomatal frequency (22.34±2.12), **facilitating efficient gas exchange, while those under 0.5% CO₂ recorded the lowest frequency (19.8±1.2)**. Chlorophyll content was likewise highest in plants cultured at 2.5% CO₂ (73.208mg/g), **correlating with increased photosynthetic efficiency, and lowest in plants at 0.5% CO₂ (56.178mg/g of leaf)**. The absence of CO₂ in the 0% treatment resulted in no detectable growth, indicating a complete lack of physiological activity.

Following the culture phase, plantlets were transferred to thermocol cups containing a 1:1:1 mixture of soil, sand, and compost for the hardening process. Plants grown at 2.5% CO₂ showed exceptional adaptation to the *Ex-Vitro* environment, achieving a survival rate of 85% and demonstrating vigorous growth. Conversely, plantlets from the 0.5% CO₂ treatment exhibited limited growth and a lower survival rate of 38%, reflecting the adverse effects of inadequate CO₂ on plantlet vigor and acclimatization.

The findings of this study emphasize the critical role of CO₂ in driving *in vitro* organogenesis, enhancing biochemical composition, and supporting robust plant growth. Among the tested concentrations, 2.5% CO₂ emerged as the most favourable, promoting superior shoot formation, metabolic activity, and adaptability to *Ex Vitro* conditions. In contrast, lower CO₂ levels (0.5% and 0%) negatively affected all parameters, including growth, nutrient assimilation, and survival rates. These results highlight the need to optimize CO₂ levels during *In Vitro* culture to maximize propagation efficiency and plantlet quality.



Graph 1



Graph 2

CONCLUSION

The research on the in vitro cultivation of *Nyctanthes arbor-tristis* L. emphasizes the essential influence of carbon dioxide (CO₂) levels on plant development, physiological traits, and biochemical properties during direct organogenesis. The findings demonstrate that elevated CO₂ concentrations significantly enhance shoot formation, nutrient assimilation, and overall plant vigor, with 2.5% CO₂ proving to be the most effective concentration. Plants grown under this condition exhibited superior growth parameters, including a higher

number of leaves, enhanced phosphorus, protein, and reducing sugar content, as well as increased stomatal frequency and chlorophyll levels, which collectively indicate improved metabolic and photosynthetic efficiency.

In contrast, suboptimal CO₂ conditions (0.5%) and the absence of CO₂ (0%) resulted in reduced growth and organogenesis, with the latter condition leading to complete failure of shoot and leaf formation. These results underscore the critical necessity of CO₂ for successful in vitro plant culture, especially under conditions designed to simulate photoautotrophic growth.



Plantlets propagated under 2.5% CO₂ also showed the highest adaptation and survival rates during the *Ex-Vitro* acclimatization phase, indicating their superior robustness and readiness for transplantation. The study confirms that optimizing CO₂ levels during in vitro culture not only enhances the efficiency of propagation protocols but also improves the quality and resilience of the plantlets.

Overall, this research provides valuable insights into the role of environmental factors in plant tissue culture and suggests further exploration of the interaction between CO₂ and other variables, such as light and humidity, to refine propagation techniques for diverse plant species.

REFERENCES

1. Rawat, Himanshi & Verma, Yashika & Saini, Neha & Negi, Neha & Chandra Pant, Hem & Mishra, Aditi & Singhal, Mayank & Khan, Arzoo & Gaurav, Naveen. (2021). *Nyctanthes arbor-tristis*: A traditional herbal plant with miraculous potential in medicine. 6. 427-440.
2. Singh, Jaspal & Singh, Amar & Singh, Ajeet. (2021). *Nyctanthes arbor-tristis*: a comprehensive review. *World Journal of Current Medical and Pharmaceutical Research*. 74-78. 10.37022/wjcmpr.v3i4.181.
3. Ashokkumar, K., Dharshini, M., Janani, T. et al. *Nyctanthes arbor-tristis* Linn. (Night Jasmine): extraction techniques, phytochemical constituents, and biological impacts of extracts and essential oil. *Futur J Pharm Sci* 10, 117 (2024). <https://doi.org/10.1186/s43094-024-00694-2>
4. Champa Rani, Sunaina Chawla, Manisha Mangal, AK Mangal, Subhash Kajla & AK Dhawan (2012), *Nyctanthes arbor-tristis* Linn. (Night Jasmine): A sacred ornamental plant with immense medicinal potentials *Indian Journal of Traditional Knowledge* Vol. 11 (3), , pp. 427-435
5. Solanki, Mansi & Rajhans, Sanjukta & Pandya, Himanshu & Mankad, Archana. (2021). *Nyctanthes arbor-tristis* Linn: a short review. *World journal of pharmacy and pharmaceutical sciences*. 10. 1047-1054. 10.20959/wjpps20213-18575.
6. Samir Panda , Deeparani Urolagin , Vijaya Kumar J , Harish N (2024) A Systematic Review Of : *Nyctanthes Arbor-Tristis*: A Conventional Herbal Remedy With Remarkable Medicinal Potential" *2IJNRD* | Volume 9, Issue 2 February 2024 | ISSN: 2456-4184 | IJNRD.ORG
7. Kozai, T. (1991). *Micropropagation under photoautotrophic conditions*. *Plant Cell, Tissue and Organ Culture*, 28, 31-39.
8. Pospisilova, J., Ticha, I., Kadlecek, P., Haisel, D., & Plzakova, S. (1999). *Acclimatization of micropropagated plants to ex vitro conditions*. *Biologia Plantarum*, 42, 481-497.
9. Deng, M. D., & Donnelly, D. J. (1993). *In vitro hardening of potato plantlets for acclimatization to harsh environments*. *Journal of Horticultural Science*, 68, 385-392.
10. Yoon, Y. J., Park, S. Y., Kim, S. H., & Paek, K. Y. (2009). *Micropropagation and ex vitro establishment of plants using photoautotrophic culture systems*. *Plant Biotechnology Reports*, 3, 235-245.
11. Solarova, J., & Pospisilova, J. (1989). *Growth and development of *Nicotiana tabacum* L. plantlets under photoautotrophic conditions*. *Acta Horticulturae*, 230, 147-154.
12. Ticha, I. (1996). *Photoautotrophic culture of plantlets: Recent trends and prospects*. *Biologia Plantarum*, 38, 13-22.
13. Vyas, S., & Purohit, S. D. (2003). *Photoautotrophic micropropagation of *Wrightia tomentosa**. *Biologia Plantarum*, 47, 125-129.
14. Joshi, M., Mehta, R., & Mathur, J. (2009). *Effect of sugar-free medium on the growth of *Chlorophytum borivilianum**. *Indian Journal of Biotechnology*, 8, 333-337.
15. Solarova, J., & Pospisilova, J. (1997). *Sucrose-free media and its effects on micropropagation*. *Plant Cell, Tissue and Organ Culture*, 51, 59-64.
16. Rahman, M. H., & Alsadon, A. A. (2007). *Influence of sucrose on potato micropropagation under photoautotrophic conditions*. *Scientia Horticulturae*, 111, 355-360.
17. Vyas, S., & Purohit, S. D. (2004). *Growth performance of *Achras zapota* under CO₂-free conditions*. *Journal of Plant Physiology*, 161, 1229-1236.